


USAGE	Biocidin® MAY BE TAKEN WITH OLIVIREX®	Biocidin® LSF MAY BE TAKEN WITH OLIVIREX®	Olivirex® MAY BE TAKEN WITH BIOCIDIN®	Biotonic™ MAY BE TAKEN WITH BIOCIDIN® & OLIVIREX®	G.I. Detox™ 1 HOUR APART FROM FOOD, SUPPLEMENTS & MEDICATIONS	Proflora® 1 HOUR APART FROM BIOCIDIN®, OLIVIREX® OR G.I. DETOX™						
BEGINNING Adults: start with the Beginning dose before increasing to Moderate or Maximum usage	1-2 drops three times a day directly on the tongue or diluted in a ¼ cup of water or juice, ½ hour before meals.  x 3 or  x 2 FOOD	Mix 1 pump with 5 oz of water. Each oz of mixture contains 1 drop of Biocidin. Start with 1 tsp of diluted mixture the first dose, hold under the tongue for 30 seconds. If no strong reaction is felt, take another tsp in 3 hours, for a total of 3 tsp the first day. The following day repeat the same way, gradually increasing the amount of mixture taken per tolerance. Work up to using the entire 5 oz of mixture in one day. Very sensitive persons use 10 oz of water, for a greater dilution.  FOOD	1 capsule twice a day.  x 2 FOOD	10 drops twice a day.  x 2	For "mop-up" of die off reactions take 1 - 2 capsules 1 hour apart from food, supplements and medications.  between meals	3 pumps into a ¼ cup of water once or twice a day, at least 1 hour apart from Biocidin®, Olivirex® or G.I. Detox™.  x 2 BIOCIDIN® OLIVIREX® G.I. DETOX™						
MODERATE Adults: gradually work up to the Moderate usage for active infections	3-5 drops three times a day, ½ hr before meals.  x 3 or  x 3 FOOD	Begin with 1 pump under the tongue per day, and gradually increase to 3 pumps a day. 	2 capsules twice a day.  x 2 FOOD	10 drops twice a day.  x 2	For general cleansing use every 2 weeks for a 3 day period. 1 - 2 capsules three times a day with a large glass of water, between meals. Take at least 1 hour apart from food, supplements and medications. 2x / Month - 3 days  x 3 FOOD	Convenient to take just upon arising and before bedtime. If you are using other probiotic products, take these with Proflora® at the same time to help improve their effectiveness. Proflora® does not require refrigeration.						
MAXIMUM Adults: if needed and well tolerated, gradually work up to the Maximum usage and maintain until symptoms improve	6-10 drops three times a day, ½ hour before meals. Maintain until symptoms improve.  x 3 or  x 3 FOOD	2 pumps three times a day.  x 3 FOOD	2 capsules three times a day in severe or acute cases until symptoms improve.  x 3 FOOD	10 drops three times a day.  x 3	<table border="1"> <tr> <td data-bbox="1438 1247 1480 1307"></td> <td data-bbox="1480 1247 2058 1307">May be taken together.</td> </tr> <tr> <td data-bbox="1438 1307 1480 1364"></td> <td data-bbox="1480 1307 2058 1364">1 hour apart from food, supplements, or medications.</td> </tr> <tr> <td data-bbox="1438 1364 1480 1421"></td> <td data-bbox="1480 1364 2058 1421">1 hour apart from Biocidin®, Olivirex® or G.I. Detox™.</td> </tr> </table> <div data-bbox="1459 1445 1575 1583">  </div> <p data-bbox="1585 1510 2016 1559">BIO-BOTANICAL RESEARCH INC.</p>			May be taken together.		1 hour apart from food, supplements, or medications.		1 hour apart from Biocidin®, Olivirex® or G.I. Detox™.
	May be taken together.											
	1 hour apart from food, supplements, or medications.											
	1 hour apart from Biocidin®, Olivirex® or G.I. Detox™.											

	Biocidin® MAY BE TAKEN WITH OLIVIREX®	Biocidin® LSF MAY BE TAKEN WITH OLIVIREX®	Olivirex® MAY BE TAKEN WITH BIOCIDIN®	Biotonic™ MAY BE TAKEN WITH BIOCIDIN® & OLIVIREX®	G.I. Detox™ 1 HOUR APART FROM FOOD, SUPPLEMENTS & MEDICATIONS	Proflora® 1 HOUR APART FROM BIOCIDIN®, OLIVIREX® OR G.I. DETOX™
USAGE MAINTENANCE Adults: for long-term immune support, after all symptoms resolve	5 drops once a day, ½ hour before meals.  x 1 /meals or 1 capsule once a day.  x 1 FOOD	1 pump once a day, 10 minutes before meals.  x 1 /meals FOOD	1 capsule once a day.  x 1 FOOD	10 drops once a day.  x 1 FOOD	Use twice a month for a 3 day period, 1-2 capsules three times a day with a large glass of water, between meals. Take at least 1 hour apart from food, supplements and medications.  x 3 FOOD	3 pumps in a ¼ cup of water once a day.  x 1 BIOCIDIN® OLIVIREX® G.I. DETOX™
CHILDREN Gradually work up to the doses listed for children less than age 12	Children under 12: 1 drop for every 10 lbs of body weight per day, in divided doses*. Take ½ hr before meals. Severe cases: up to 2 drops per 10 lbs per day may be used if well tolerated. Maintenance: ½ drop per 10 lbs per day may be used for long-term immune support. *Divided doses: Divide the dose to twice a day (a 40 lb child could take 2 drops AM and 2 drops PM, for a total of 4 drops a day.) 1 - 2  / 10lbs x 2 FOOD	Dilute 1 pump into 5 oz of water. Each oz of mixture contains 1 drop of Biocidin. Start with 1 tsp of diluted mixture, per 10 lbs of body weight for the first dose, hold under the tongue for 30 seconds. If no strong reaction is felt, take the same dosage in 3 hours, for a total of 3 dosages the first day. Work up to 2 Tbsp per 10 lbs of body weight per day.* Take 10 minutes before meals. Severe cases: Up to 4 Tbsp of mixture per 10 lbs of body weight per day if tolerated. *Divided doses: divide the dose to twice a day. A 40 lb child could take 4 Tbsp morning and 4 Tbsp evening, for a total of 8 Tbsp/day. 1 Tbsp / 10lbs x 2 FOOD	Children ages 7 -12 that can swallow a capsule may use ½ capsule twice a day.  x 2 Teenagers may use 1 capsule twice a day.  x 2 FOOD	Children ages 7 -12: 5 drop three times a day.  x 3 Teenagers may use 10 drops twice a day.  x 2 FOOD	Children under 50 lbs: Use twice a month for a 3 day period. 1/2 capsule twice a day. Eat a diet high in leafy greens and low in refined carbohydrates. May be mixed with juice or other liquids. 2x / Month - 3 days  x 2 FOOD	Do not use Proflora® with children under 2. 2 pumps in a ¼ cup of water twice a day. Best taken upon rising and at bedtime. Proflora® does not require refrigeration.  x 2 BIOCIDIN® OLIVIREX® G.I. DETOX™
PREGNANCY AND NURSING MOTHERS	Do not use during pregnancy. Breast feeding mothers may use at ½ the recommended adult dose if well tolerated.	Do not use during pregnancy. Breast feeding mothers may use at ½ the recommended adult dose if well tolerated.	Do not use during pregnancy. Breast feeding mothers may use at ½ the recommended adult dose if well tolerated.	Not recommended for use during pregnancy.	Not recommended for use during pregnancy.	Do not use during pregnancy. May be used for breast feeding mothers.
SPECIAL PRECAUTIONS	Do not use before surgery as some herbs may have a blood thinning effect. Discontinue use should adverse reactions occur.	Do not use before surgery as some herbs may have a blood thinning effect. Discontinue use should adverse reactions occur.	Do not use before surgery as some herbs may have a blood thinning effect. Discontinue use should adverse reactions occur.	Discontinue use should adverse reactions occur.	Discontinue use should adverse reactions occur.	Discontinue use should adverse reactions occur.